

MICROBIAL DIVERSITY AND ITS ROLE IN ENHANCING SOIL FERTILITY: A COMPREHENSIVE REVIEW

Asiya Akbar¹, Dr Saman Zahra^{*2}, Qasid Hussain³, Usman Waheed⁴, Muhammad Rehman⁵

¹Department of Agriculture and Environmental Sciences, The Islamia University of Bahawalpur.

²Department of Biology, University of Okara, Pakistan

³Scientific Officer, Agricultural Research Institute Jaffarabad BARDC PARC

⁴Faculty of Crop Production, Sindh Agriculture University Tandojam Hyderabad Sindh.

⁵Department of Environmental Science, Faculty of Basic and Applied Sciences, International Islamic University, Islamabad

¹asiyaakbar413@gmail.com, ²saman.zahra@uo.edu.pk, ³qasidh@gmail.com,

⁴muhammadrehmanburdibaloch@gmail.com, ⁵osman.waheed@outlook.com

DOI: <https://doi.org/10.5281/zenodo.14851565>

Article History

Received on 23 December 2024

Accepted on 23 January 2025

Published on 10 February 2025

Copyright @Author

Corresponding Author: *

Dr. Saman Zahra

Department of Biology, University of Okara, Pakistan

saman.zahra@uo.edu.pk

Abstract

Soil microbial diversity is of utmost significance in improving and maintaining soil fertility, a prerequisite for sustainable agriculture. Microorganisms-bacteria, fungi, actinomycetes, archaea, and protozoa, perform all soil functions such as nutrient cycling, organic matter decomposition, and soil aggregation and, directly or indirectly, provide an essential nitrogen fixation, solubilization of phosphorus, and mobilization of potassium for the plants. They improve soil structure, moisture retention, pH level regulation, and root growth of nitrogen-fixing rhizobia and mycorrhizal fungi. However, microbial diversity is influenced by several geological factors, namely texture, pH, organic matter, moisture, agricultural practices, and temperatures. Conservation and Sustainable farming practices understand the microbial diversity contribution as organic farming, less tillage, and cropping system diversification support them, whereas normal practices usually reduce it. Modern methodologies like sequenced 16S rRNA, metagenomes, and metatranscriptomes have made a revolutionary change regarding soil microbial community studies for their deep and detailed insights into the structure and function of these parts. Understanding microbial diversity and its interactions helps formulate better soil management strategies. This review emphasizes the importance of microbial diversity for enhancing soil fertility while discussing sustainable agricultural practices in support of maintaining a healthy soil ecosystem.

INTRODUCTION

Importance of Soil Fertility in Agriculture

Its very essence hinges on agricultural productivity and sustainability, such as sufficiency: the quality of the soil to be efficient in supplying the nutrients at once, and in sufficient amounts to the plants, at the

time of growth and development, is called soil fertility. A fertile soil will give rise to healthy crops and boost yields and food security alongside economic stability in agricultural systems. For millennia, human civilizations have relied on the

fertility of soils for survival and welfare; these often serve as the foundation of powerful agrarian societies[1, 2].

Nevertheless, excessive tillage, monoculture cropping practices, and overuse of chemical fertilizers and pesticides, modern agricultural practices have worsened soil fertility declines in many regions. The consequences of declining soil health include reduced productivity by crops, loss of biodiversity, and increased susceptibility to environmental stresses such as drought and flood damage. Thus, understanding and improving soil fertility have become global priorities to satisfy demands for feeding an ever-increasing population and addressing the effects of climate change[3].

Role of Microbial Diversity in Soil Health and Fertility

Soil microbial diversity has a very wide meaning and classical understanding. It refers to the variety of all kinds of microorganisms in a soil ecosystem—bacteria, fungi, archaea, protozoa, and algae. Just looking at the soil, however, is not sufficient evidence that microorganisms form a big part of the overall organismic world. If perceived, most people regard soil as simply a lifeless substrate on which plants grow. Rather than that, it is a dynamically living system in which billions of cells live, thrumming with life. For almost any amount in a gram of soil, you can find billions of microbial cells representing thousands of species. They constitute a whole lot of important functions that directly concern soil properties, such as their physical, chemical, or biological elements[4].

Soil microbes are the requisites for nutrient cycling. The microorganisms facilitate organic matter decomposition into simpler compounds that release nutrients important for growth such as nitrogen, phosphorus, and potassium. For example, nitrogen-fixing bacteria like *Rhizobium* and free-living diazotrophic microbes such as *Azotobacter* transform atmospheric nitrogen into forms absorbable and usable by plants. The function of phosphate-solubilizing bacteria and fungi consists of mobilizing the phosphorus in adhesion with insoluble minerals and making it available to plants [5].

Instead of nutrient cycling, soil microbes enhance the soil structure by the formation of biofilms and

exopolysaccharides for binding soil particles into aggregates that hold water and stand aeration. Fungi such as mycorrhizal fungi stretch their hyphae into the soil, producing networks that enhance both soil and nutrient uptake in plants. Microbial activity also regulates soil pH, detoxifies polluting chemicals, and suppresses most soil-borne pathogens, which enhances further plant welfare[6].

Objectives and Scope of the Review

The potential microbial diversity in soil fertility has drawn great attention from scientists, policymakers, and farmers. Questions are being raised about traditional agricultural practices for their environmental consequences, yet there is a call for the adoption of microbial soil abilities in management strategies to create environmentally friendly soil systems. This paper aims to comprehensively understand the relationships between soil microbial diversity and fertility mechanisms through which microorganisms have been beneficial to soil health and productivity[7].

The specific objectives of this review are as follows:

Be aware of the diversity of soil microorganisms visible from various functional roles among bacteria, fungi, archaea, and many others. They are responsible for different nutrient cycles and the decomposition of organic materials and structures inside soil. Evaluate how various environmental and anthropogenic factors would influence microbial diversity and soil fertility. Examine different contemporary approaches to the measurement of microbial diversity and how such techniques have been used most widely in soil science. Understand microbial diversity's importance for sustainable agriculture, and its response to world challenges in food security and climate change.

This is to say that the review will systematically pursue each of these objectives. It starts with an introduction to the subject of soil microbial diversity and its main components: an examination of the role microorganisms play in improving soil fertility. This is then followed by reviews of the interaction between soil microbes and plants, environmental factors affecting microbial diversity, and current advancements in techniques of microbial research. Further, the review considers the application and management of soil about microbial diversity, while

the conclusion indicates the gaps in the knowledge and future research.

In the end, the review highlighted the very fact that microbial diversity is indeed valuable as a current natural resource in maintaining soil health and agricultural productivity. With the world, increasing, challenges are piling such as population increases, land degradation, and climate change; conserving and using soil microbial diversity for the productive keeping of resilient and sustainable agricultural systems becomes one of the promising ways forward.

Understanding soil microbial diversity

Definition and Components of Soil Microbial Diversity

Microbial diversity in soil comprises different species and population numbers of microorganisms existing in the soil ecosystem. This diversity involves different forms of microbial species, genetic variation among those species, and differing functional roles these microorganisms execute in the soil environment. Microbial diversity is very important to soil health in that it carries out specific vital processes such as nutrient cycling, organic matter decomposition, and regulation of soil-borne pathogens[8].

The soil microbial community is constantly in turmoil since organisms interact with one another and with the environment in different ways. These interactions of the soil ecosystem contribute to its resilience and productivity. Microbial diversity can be subdivided into three broad dimensions: the number, and variety of microbial species present in the soil; genetic variation within and between microbial populations; biochemical functions, and the ecological range of functions performed by soil microorganisms.

Each of these dimensions plays a substantial role in maintaining soil fertility and ensuring the sustainability of agricultural systems. High microbial diversity often signifies a healthy and robust soil ecosystem capable of adapting to environmental stresses such as drought or contamination[9].

Methods:

We concluded this literature by using digital platforms like Google Scholar and Pubmed. The following terms were Increasing soil fertility as a factor in the sustainability of agriculture and resilience to climate change, Soil fertility in arid

lands: strategies for sustainable management and fertilization, in Sustainable Agriculture under Drought Stress. Linking soil microbial diversity to modern agriculture practices: a review. Soil microbial diversity and the sustainability of agricultural soils The Role of Sustainable Farming Practices in Enhancing Soil Fertility and Crop Yield. Dialogue Social Science Review, Diversity, functions, and stress responses of soil microorganisms. Plant Microbiome: Stress Response, All literature reviews, original article papers and all cases, and reports regarding aspects. The cross references from all these papers and case reports were also included.

Types of Soil Microorganisms

Soil is home to a vast array of microorganisms, each with unique roles and contributions to the soil ecosystem. The major types of soil microorganisms include bacteria, fungi, archaea, actinomycetes, protozoa, and algae[10].

1. Bacteria

Soil comprises by far the most numerous and diverse group of microorganisms: bacteria. They are unicellular, prokaryotic organisms that carry out a variety of functions necessary for soil health. Thus, bacteria participate in nitrogen fixation, nitrification, and denitrification alongside work. For example, such nitrogen-fixing bacteria as Rhizobium and Azotobacter will change atmospheric nitrogen into ammonia, a form that plants can uptake. Saprophytic bacteria will break down organic matter and release nutrients to help form humus. Plant Growth-Promoting Rhizobacteria (PGPR) are a specific group of bacteria that enhance plant growth by producing phytohormones, solubilizing nutrients, and suppressing pathogens [6, 11]. Pseudomonas and Bacillus are instances. Certain bacteria do degrade environmental pollutants while improving soil quality.

Bacteria seem to have very wide adaptability. They thrive in soil conditions ranging from acidic ones to alkaline ones and nutrient-rich ones to nutrient-poor ones.

Fungi

Fungi are eukaryotic organisms that play a major role in the soil ecosystem. They have been classified based on their ecological activities into three groups.

Two major groups of fungi complete the decomposition process: they break complex organic materials such as cellulose and lignin into simpler ones. Examples of such fungi include *Trichoderma* and *Penicillium*. These are symbionts that form a mutualistic relationship with plant roots, enhancing nutrient (mainly phosphorous) absorption and improving the resistance of plants to different environmental stresses. Among many examples of Arbuscular mycorrhizal fungi (AMF), some are specific such as *Fusarium* and *Phytophthora* are plant pathogens causing diseases in crops.

Fungi are important in soil aggregation and stabilization by forming networks of hyphae binding soil particles and producing glomalin, a glycoprotein that improves soil aggregation and stability[12].

3. Archaea

Archaea, differently cognized to bacteria, are prokaryotes that have a distinct genomic and functional configuration. These organisms are associated with extreme environmental conditions such as highly saline, acidic, or high-temperature environments; however, most of the time they are just soil organisms. Methanogen archaea produce methanol while methanotrophic archaea oxidize it; they thereby participate in the carbon cycle. Ammonia oxidizing archaea (AOA) also contribute to nitrification; their principal activity, however. That has been recorded in nutrient-poor or acidic soils[13].

4. Actinomycetes

Actinomycetes, what are they? Yes, they are filamentous bacteria, which show properties of both bacteria and fungi. They are most abundant in soils that consist of high organic mass and contribute to the typical "earthy" smell of soils. Actinomycetes degrade complex organic substances into simple ones, such as chitin and cellulose. Many actinomycetes are antibiotic producers, such as *Streptomyces*, contributing to their efficacies against soil-borne pathogens toward the enhancement of plant health. Actinomycetes exist for nitrogen and phosphorus cycling, thereby making these nutrients available for plant uptake. Actinomycetes are the most active during composting and in soils with low pH [14].

5. Protozoa and Algae

- Protozoa: Protozoa are unicellular eukaryotic organisms that feed on bacteria, fungi, and other soil microorganisms. Grazing on bacteria, protozoa help regulate microbial populations and release nutrients available in plant-usable forms, such as nitrogen. They also improve soil structure by mucilage, binding soil particles to one another[15].

- Algae: Photosynthetic organisms are called algae, and they play a significant role in soil fertility, especially for the upper layers of soil. They are added to soil mostly through the activities of cyanobacteria, which represent the blue-green algae. They are particularly important in soils that are dry and semi-dry; usually fix atmospheric nitrogen and form biological soil crusts, thus minimizing erosion while improving the water-holding capacity of the soil[16]. Although protozoa and algae are less abundant than bacteria and fungi, they play critical roles in maintaining soil ecosystem balance.

Factors Influencing Microbial Diversity in Soils

Soil microbial diversity is influenced by a wide range of environmental, biological, and anthropogenic factors. These factors determine the composition, abundance, and activity of soil microorganisms.

1. Soil Properties:

Poor soil limits the types of organisms that can grow in it. This is particularly true for pH: fungi are better able to thrive in acidic soils, while there will be more different kinds of bacteria in neutral to alkaline soils. It will be sandy soils that produce a rather limited microflora since they are unable to retain water well. Better conditions for microbes are clay soils. High organic matter content supports microbial diversity by providing energy and nutrients[17].

2. Climate:

High temperatures as well as moisture affect microbe activities and diversity. For instance, activity levels in bacteria are higher than usual in warm and moist conditions while fungi predominate in cool, dry conditions[18].

3. Role of Microorganisms in Soil Fertility

Microorganisms are crucial for the maintenance and improvement of soil fertility. Microorganisms

perform numerous biochemical processes that enhance the availability of nutrients, decompose organic matter, induce the soil structure, and stabilize the environment of the soil. These processes are the basis of healthy and sustainable agricultural systems[19].

Nutrient Cycling and Availability

Microorganisms are very important in cycling and mobilizing vital nutrients so that they can be available to plants within the soil. Some of the major activities offered by microorganisms include:

Nitrogen Fixation

Nitrogen is an essential plant nutrient. Most plants, however, cannot directly utilize atmospheric nitrogen (N_2). Nitrogen-fixing microorganisms convert atmospheric nitrogen into ammonia (NH_3), which can, in turn, be utilized by the plants, through biological nitrogen fixation. An example is that of Rhizobium bacteria that form nodules on the roots of leguminous plants and fix nitrogen in return for carbon compounds. Nitrogen fixation is also contributed to in the rhizosphere by organisms such as Azotobacter and Clostridium. Some bacteria such as Azospirillum change the availability of nitrogen by associating closely with the roots of non-leguminous plants.

Thereby reducing the need for chemical nitrogen fertilizers, these mechanisms further enhance sustainable agriculture through nitrogen-fixing organisms [20].

Phosphorus Solubilization

Apart from nitrogen, phosphorus is another essential nutrient for plants, but it cannot be easily available for plants as it forms insoluble compounds in soil. Phosphate-solubilizing microorganisms (PSMs) produce organic acids, enzymes, and chelating agents that make phosphates soluble to convert this insoluble phosphorus into available forms for plants. The genera like Pseudomonas, Bacillus, and Rhizobium are commonly known as effective phosphate solubilizers. Some species, such as Aspergillus and Penicillium, solubilize phosphorus effectively. [21].

Straight from PSM, phosphorus uptake by plants can be promoted, leading to improved root and overall plant health.

Mobilization of Potassium

Potassium is a macronutrient which is needed by plants for their metabolism. Much of the potassium occurs in mineral forms that are not soluble. Potassium solubilizing microorganisms (KSMs) solubilizing or releasing potassium from mineral sources such as feldspar and mica through organic acids and enzymes. Bacillus mucilaginosus and Paenibacillus have been potassium-solubilizing bacteria. Such fungi species can also be potassium-releasing. Such activities can help an organism by better nutrition, increased productivity, and resistance to the abiotic stresses for the crops[22].

Organic Matter Decomposition and Humus Formation

The decay of organic matter is a process and activity involving soil microorganisms, which break down plant residues, animal excreta, and other organic matter into simpler compounds. Decomposition has several benefits. Firstly, it releases certain nutrients such as nitrogen, phosphorus, and sulfur and makes them available for plant uptake. The second benefit includes humus, a stable organic matter fraction that increases the fertility of the soil. Microorganisms including actinomycetes and fungi play an important role in humus formation due to their ability to decompose such complex compounds as cellulose, lignin, and chitin[23].

There are examples like Trichoderma and Penicillium, which have the cellulose and lignin-decomposing ability of organic matter along with examples like proteolytic, saccharolytic, and lipolytic microorganisms. Some other specific examples include Pseudomonas and Bacillus. They are key organisms involved in the decomposition of complex organic compounds such as cellulose and lignin, thus assisting in humus generation. Nutrient recycling, improved soil structure, and water-holding capacity are other benefits of decomposition[24].

Improvement of Soil Structure and Aggregation

Microorganisms have a very important task to accomplish in soil structure and aggregation, which

is vital for aeration, infiltration of water, as well as root growth. Bacteria secrete exopolysaccharides through which soil particles are bound to form an aggregate of stable structures in continuity. This further improves porosity in the soil and prevents erosion. Mycorrhizal fungi have been shown to form extensive networks of hyphae that can bind together soil particles. This has been shown that they produce glomalin-a protein enhancing aggregation. Earthworms are not microorganisms but rely on microbial activity on organic matter. This also contributes to the formation of soil aggregates, from casting[25].

Soil aggregation improvement enhanced water retention, decreased compaction, and increased strength in the plant root systems. Microbial Contributions to Soil pH Regulation Microbial activity and nutrient availability have been largely affected by soil pH. Microbes through the organic acids, ammonia, and other organic compounds are

actively involved in the regulation of the soil pH. Microbial or phosphate-solubilizing bacteria and fungi secrete organic acids that lower soil pH by mobilizing such nutrients into bioavailable forms, including phosphorus and potassium. Ammonia-producing bacteria and archaea may raise the soil pH through alkaline compounds released during nitrogen transformation. Microorganisms take part in the buffering capacity of soils by carbonates and bicarbonates cycling and thus stabilizing fluxes in pH[26].

Microorganisms decipher soil pH into a state where they invoke processes of nitrification, e.g. Nitrosomonas and Nitrobacter. On the other hand, some fungi like Aspergillus have acids secreted into the environment that dissolve calcium phosphates, leading to a reduced pH in areas on the soil. It is through these mechanisms that microorganisms maintain an optimal range for plant growth and nutrient availability.

Microbial Role	Key Processes	Examples of Microorganisms	Benefits	References
Nutrient Cycling	Nitrogen Fixation	Rhizobium, Azotobacter, Clostridium	Converts atmospheric nitrogen to ammonia.	[26]
	Phosphorus Solubilization	Pseudomonas, Bacillus, Aspergillus	Solubilizes insoluble phosphates for plants.	[26]
	Potassium Mobilization	Bacillus mucilaginosus, Paenibacillus	Releases potassium from mineral sources.	[25]
Organic Matter Decomposition	Breakdown of organic residues	Trichoderma, Penicillium, Bacillus	Releases nutrients and forms a stable humus.	[27]
Soil Structure Improvement	Soil aggregation and stability	Mycorrhizal fungi, Pseudomonas	Improves aeration, water retention, and roots.	[28]
pH Regulation	Organic acid/ammonia production	Nitrosomonas, Aspergillus, Nitrobacter	Stabilizes pH for optimal nutrient uptake.	[27]

Table: The critical roles and microorganisms involved

Interactions Between Microorganisms and Plants

There is a dynamic interaction between soil microorganisms and the root systems of plants, most of which are related to plant growth, development, and health. These interactions could be mutualism or plant growth promotion and protection from environmental stress and pathogens. Understanding these interactions is, therefore, important to realize the microbial potential for sustainable agriculture advancement [29].

These interactions represent examples of mutualism between microorganisms and plants: in this case,

microorganisms help plants obtain essential nutrients, while the latter benefit from the nutrients supplied by the microorganisms and their protective mechanisms. Such microbes include rhizobia, which are soil bacteria involving legumes in symbiotic association. They inhabit root nodules fixed into plant ammonia because of nitrogen captured from the atmosphere. They invade root hairs, lead to nodule formation, and fix nitrogen using nitrogenase enzymes. This has enhanced nitrogen availability while reducing dependence on chemical fertilizers.

Foreign examples include *Rhizobium* species associated with beans, peas, and soybeans [30].

Mycorrhizal fungi are involved in making important mutualistic relationships with plant roots, which improve nutrient and water uptake and protect soil pathogens. Arbuscular mycorrhizal fungi penetrate the root cortical cells and exchange nutrients, while ectomycorrhizal fungi form an external fungal sheath around roots. These fungi increase phosphorus, nitrogen, and micronutrient absorption; increase drought tolerance; and improvements in resistance to soil pathogens. Examples include AMF species like *Glomus* and ectomycorrhizal fungi like *Pisolithus* associated with forest trees [31, 32].

Plant growth-promoting rhizobacteria (PGPR) are free-living bacteria that associate with plant roots and improve their growth through a range of mechanisms. These include enhancing nutrient uptake, stress tolerance, and pathogen resistance. Some PGPR can solubilize phosphorus, mobilize potassium, chelate iron through producing siderophores, and make such nutrients available for plants; produce hormones like auxins, gibberellins, and cytokinins to stimulate root and shoot development; and trigger defensive responses in plants, resulting in enhanced plant immunity, also known as induced systemic resistance. Predominant examples of this include *Pseudomonas fluorescens*, *Bacillus subtilis*, and *Azospirillum*. It increases root growth, yield improvement, abiotic stress mitigation, and decreases chemical application requirements [33].

Endophytic microorganisms live inside plant tissues without causing harm. They establish a symbiotic relationship and provide benefits like enhanced nutrient acquisition, growth promotion, and stress resistance. Endophytes mobilize nutrients and directly deliver them to plant tissues. They bring plants durability against the adverse conditions of drought, salinity, and heavy metals, by inducing compounds like proline, which are known as stress-related compounds. Endophytic bacteria and fungi produce many anti-microbial agents or compete for resources against pathogens to decrease disease incidence. Such examples include endophytic bacteria, *Burkholderia*, and *Enterobacter*, with the skilled fungal examples being *Fusarium* and


Trichoderma. These microorganisms make more tolerance to the plants and, thus, improve crop productivity [34].

Microbial interactions are the core of plant health and productivity. From nutrient acquisition to growth promotion, stress tolerance, and protection from pathogens, these relationships stress the point that microbial diversity must be tapped into for the sustainability of agricultural practices. An in-depth understanding of these relationships could probably lead to cutting-edge innovations in enhancing soil health, thereby reducing chemical input dependency and ensuring food security [35].

Impact of Environmental Factors on Microbial Diversity

Environmental factors play an essential role in determining the microbial soil diversity. They determine the number, activities, and types of microorganisms living in the soil and then influence the soil fertility and plant health. Soil texture, pH, organic matter content, temperature, moisture changes, and agricultural activities are all important factors that define the ecology of a soil ecosystem.

The key factors affecting microbial diversity are soil texture, pH, and organic matter content. Soil texture which is the amount of sand clay and silt particles determines the amount of air and moisture kept in soils plus nutrient allocation [17]. Organic matter-rich loamy soils are more likely to promote high microbial diversity because they are good conditions for microbial growth and nutrient cycling. On the other hand, heavy clay soils usually restrict the activity of microorganisms due to their poor aeration, while sandy soils may also restrict the availability of nutrients. Soil pH further affects the distribution of microbes. The microorganisms are adapted to different pH ranges. A few examples; the *Azotobacter* and *Bacillus* bacteria mostly thrive in neutral to slightly alkaline soils, while fungal members such as *Penicillium* show a greater preference for acidic conditions. Extreme acidic or alkaline pH conditions reduce the diversity and activity of microbes by limiting the cycling of nutrients and the fertility of soils [36].



Bar chart showing the positive and negative impacts of environmental factors on microbial diversity.

The amount of organic matter present in soil decayed by dead plants, animals, and microbes forms the major source of energy and nutrients for soil microbes. Generally, organic matter at higher levels supports a diversified and more active microbial population and provides enhanced effectiveness in nutrient cycling, soil aggregation, and overall soil health, while depletion of organic matter due to poor management of land reduces microbial diversity and nutrients in soil [37].

Soil temperature and moisture vary significantly among countries, and are among the most important environmental factors that influence microbial diversity. Microorganisms are affected by temperature and usually have a direct relationship between growth and surrounding temperature conditions. Biologically, warm active months are said to favor the proliferation of soil microorganisms at higher soil temperatures, as typically found in temperate countries. However, too high a temperature may cause stress to soil microbial communities and decline in that abundance. In the same way, barely freezing temperatures slow microbial activities per se but may allow the survival of cold-tolerant species [38].

Moisture is one of the critical factors that determines microbial activities. Soil moisture has an influence on the availability of water to microbial processes; variations in moisture may have a direct effect on the ability of microorganisms to survive and the diversity of their populations to dry conditions will dry out microorganisms, while too wet conditions will limit oxygen supply and tend to promote an anaerobic situation. Moderately moist conditions would be expected to favor a relatively wider range of microorganisms and therefore help support a more balanced and diverse microbial community [38].

Changes in agricultural methods and conversion of land from one use to another have a very great effect on microbial diversity. Practices like mono-crop cultivation, the addition of heavy chemical fertilizers and pesticides, and intense tillage activity are thought to drastically reduce the microbial diversity available and destroy its natural balance within soil ecosystems. Intriguingly, even when widely practiced, synthetic fertilizers can change soil pH and nutrient availability, thus favoring certain groups of microbes at the expense of others. In addition, the use and overuse of pesticides may also inhibit beneficial soil microorganisms that perform important functions

such as nitrogen fixation and organic matter decomposition [39].

This would be one means by which microhabitats change as time goes by, in terms of land-use changes like deforestation, urbanization, and soil erosion. Deforestation would cause the following: reduced organic matter input disrupted nutrient cycling, and decreased microbiological diversity. All three phenomena-induced soil compaction, reduced organic matter, and increases of pollutants in soils may cause stress to microbial populations. Agricultural measures that would ensure minimum such impacts promote healthy microbial community and increased fertility including crop rotation, conservation tillage, and organic farming [40].

Because soil texture, pH, organic matter, temperature, moisture, and farm practices all play significant roles in the microbial diversity within soils. Very important factors responsible for influencing nutrient cycling, maintaining soil health, and furthering the goals of sustainable agriculture are the understanding of such environmental factors and their effects on microbial communities. Optimizing these environmental factors is an important aspect of improving fertility in soils while contributing to sustainable food production systems [41].

Modern Techniques for Assessing Microbial Diversity

Modern techniques for assessing microbial diversity have evolved significantly, incorporating both conventional cultivation methods and advanced molecular approaches. They make it possible for researchers to strive for a wider understanding of microbial communities inhabiting different environments, including soils. Cultivation methods reveal specific microorganisms that can grow under controlled laboratory conditions, while molecular methodologies give a fuller view of the entire microbial community, including the unculturable ones [42].

Traditionally, cultivation-based techniques have been utilized for detecting microorganisms present in a sample by isolating them from the environment and growing them in nutrient-rich media such techniques have opened up an entirely new world in understanding microbial diversity, but they are

limited by the inability of many microorganisms, particularly those with specialized or slow growth requirements, to be cultivated in the laboratory. Hence, it may more often describe an environmental diversity[43]. Nonetheless, despite these limits, cultivation will still serve the purposes of studying particular organisms, describing new species, and studying microbial physiology. Selective media can be used for specific groups of microorganisms (e.g., bacteria, fungi, actinomycetes) to investigate their role in different ecosystems.

Molecular techniques have impacted the study of microbial diversity, as they allow scientists to analyze, directly from environmental samples, the DNA or RNA present in these and to skip culture. This allows the detection and identification of an even larger range of microorganisms, including those difficult or impossible to culture [44].

The application of 16S rRNA sequencing has become one of the most prominent molecular approaches for the identification of bacteria and archaea in environmental samples. This ribosomal RNA gene is conserved across all species of bacteria and archaea, although some regions are variable and specific to different groups. By amplification and sequencing of the 16S rRNA gene from environmental DNA, it is possible to identify and classify microorganisms according to their genetic sequences. With this method, potential knowledge about the microbial diversity of ecosystems such as soil, lakes, and human microbiomes has increased greatly. In addition, this method allows for culturing of unculturable microorganisms and gives an idea about the structures of their communities [45].

Unlike the earlier microbiological method of isolating individual microorganisms, metagenomics includes sequencing all genetic material present in environmental samples. This comprehensive method provides not only a complete overview of the entire microbial community consisting of bacteria, archaea, fungi, viruses, and other microorganisms but also allows for interpretation of the microbes within specific environmental samples.

This means that metagenomic sequencing provides knowledge about the recognized one's functions (potential contact functions) by collecting and studying those with different known and newly discovered microbes according to community-mosaic

genome assembly. Moreover, metagenomics enables one to understand microbial diversity across ecosystems by comparing metagenomic data from various environments and discovering new metabolic pathways that would otherwise remain buried under traditional approaches to microbial interactions [46]. It is metatranscriptomics that goes alongside metagenomics with the difference of being RNA-centric finds its miniscule focus on analyzing environmental samples for DNA. Since RNA points towards the dynamic activity of microorganisms, through metatranscriptomics, one can study microbial communities and how and which genes tend to be expressed at a given time. It brings conceptualization around the functional roles of microorganisms in an environment and how microbial communities respond to environmental changes. Linkages between gene expression patterns the microbial functions provide clarity on nutrient cycling, stress responses, and pathogenicity [47].

Functional analysis and bioinformatics tools play a crucial role in the interpretation of complex data generated by molecular techniques. Functional analysis allows researchers to assess roles and metabolic potential in the microbial community: degrading organic matter, fixing nitrogen, or producing bioactive compounds. In the case of functional genes directly related to nitrogen fixation, carbon cycling, and antibiotic production, the genetic sequences will model their functional diversity [46].

Processing and analyzing huge data generated from sequencing technologies is made possible by bioinformatics tools. With the help of these tools, one can perform sequence alignment, taxonomic classification, and functional annotation to figure out microorganisms and their possible functions in processes. Some popular computational biology tools for microbial diversity studies are QIIME (Quantitative Insights Into Microbial Ecology), Mothur, Kraken, and others. These tools can integrate data from different sequencing technologies and allow researchers to perform comparative analysis and provide images of the microbial community structure and functioning [48].

The microbial diversity assessment modern techniques are significantly improving an understanding of environmental microbial

communities. Some cultivation-based methods still offer the possibility of isolation and characterization of target microorganisms, whereas molecular techniques such as 16S rRNA sequencing, metagenomics, and metatranscriptomic analysis provide more exhaustive insights into both the diversity and functionality of microbial communities. Functional analyses together with bioinformatics tools are indispensable for interpreting any data obtained from these molecular techniques. Therefore, with these techniques, it becomes possible to conduct effective and accurate measurements of microbial diversity, which are very useful for various applications including agriculture, ecology, medicine, and biotechnology [49].

Microbial Diversity and Sustainable Soil Management

Microbial diversity is a key factor in the health and fertility of soils and sustainable practices of land management. It is an important aspect in the context of sustainable soil management for the long-term productivity of soils, nutrient cycling, and reduction in chemical inputs. Healthy microbial communities in soils can be developed through practices such as organic farming; the use of microbial inoculants, biofertilizers, biostimulants, and composting. These not only encourage microbial diversity but work towards an overall sustainable agriculture [5].

Microbial inoculants and organic farming are two pillars of sustainable soil management. Organic farming encourages soil health without the use of synthetic pesticides and fertilizers that harm soil microbial communities. The use of natural inputs promotes a diverse and active microbial population. One of the cornerstones of organic farming is soil organic matter, an important source of food and habitat for a variety of soil microorganisms, enhancing soil structure, nutrient cycling, and ultimately the health of the plants [50].

Microbial inoculants are effective microorganisms applied to soil for the enhancement of specific microbial functions. There is a range of inoculant organisms, which can be bacteria fungi, or other organisms that stimulate plant growth or supply nutrients into the soil. An example is rhizobial inoculants to improve nitrogen fixation of leguminous crops, while mycorrhizal inoculants

improve nutrient uptake attained through symbiotic relationships with plant roots. It will help restore and/or enrich microbial diversity that has been depleted with conventional farming practices—the introduction of beneficial microorganisms through inoculants. These types of microbial inoculants also have the potential to improve resistance to plant diseases, soil compaction, and soil fertility important aspects of sustainable agriculture [51].

Yet another way of improving soil microbial diversity is through the use of biofertilizers and biostimulants. Biofertilizers are living organisms introduced into soil or plants to enhance soil fertility. Such living organisms contribute benefits through nitrogen fixation, organic matter breakdown, and making nutrients such as phosphorus more available to plants. Under such situations, the need for chemical fertilizers is lowered while promoting the growth of beneficial microbes that build larger and more diverse communities. Some examples of biofertilizers include nitrogen fixers like *Rhizobium* and *Azotobacter*, and phosphorus solubilizers such as *Bacillus* and *Pseudomonas* [51].

On the contrary, biostimulants are organic or living compounds that fortify plant growth and health through natural action in plants and soil. These agents are quite different from fertilizers because they do not directly supply plant nutrients but influence the plant to take up nutrients, enhance nutrient availability through stress tolerance, and/or encourage microbial activity in the soil. From seaweed extracts, humic substances, or even beneficial microorganisms, biostimulants can be derived from a variety of natural sources. Notably, biostimulants could help improve soil microbial diversity and increasingly resilient, more productive agricultural systems through plant-microbe interactions to improve microbial activity [9].

Hence, composting and its effect on microbial diversity have been considered as one of the significant practices for sustainable soil management. Generally speaking, composting means the controlled decomposition of organic residues, manure, food wastes, etc., to result in a nutrient-rich humus for further use in soil health improvement. The composting process is mediated by a large variety of microorganisms - bacteria, fungi, and actinomycetes, which break up organic matter into

smaller, simpler compounds. This microbial diversity involved in composting is essential for the efficient breakdown of organic matter and the production of stable humus [52].

Microbial diversity can be very much increased by the addition of compost to soil because of the wide variety of microorganisms, many of which are beneficial for plant health. The addition of compost increases the number of microbes that are capable of decomposing organic materials, recycling nutrients, and improving soil structure. Furthermore, the diverse microbial communities found in composted materials also contribute to disease suppression, nutrient cycling, and the overall resilience of soil ecosystems. Apart from adding to the marvelous variety of soil microbes, the application of compost also benefits soil concerning fertility and sustainability over time. [53].

Microbial diversity is important for the practices of sustainable soil management. These include organic farming practices, microbial inoculants, biofertilizers, biostimulants, and composting, which help maintain or build up microbial diversity in soils. Such benefits also include soil fertility improvement, nutrient cycling, and increased plant resilience by reducing the use of chemical inputs. Thus, healthy microbial communities have been established for the long-term viability of agricultural systems and, ultimately, for environmental health contributions [54].

Discussion

Microbial diversity is one of the most important aspects of soil health, which regulates several soil functions and processes requisite for a sustainable agricultural system. Microorganisms, including bacteria, fungi, actinomycetes, archaea, protozoa, and algae, form a wide variety of their kind. These microorganisms are involved in nutrition cycling, organic matter decomposition, disease suppression, and overall soil fertility. The understanding of microbial diversity and its dynamics is paramount to formulating better strategies for enhancing soil health and conservation in sustainable farming practices [55].

Factors such as soil texture, pH, organic matter content, temperature, and moisture level affect the diversity of microorganisms in the soil. These parameters govern the kinds of microorganisms

adapted to biotic terrestrial features. For instance, soil pH is a measure of the sorts of microbial communities that can thrive therein; certain acidic soils have favored some groups while alkaline have been over others. Organic matter content is a reservoir of energy and nutrition for soil microorganisms, which ensures that very diverse populations exist. Soil texture-specified drainage capacity influences the availability of water and oxygen to microbes and controls their activity. [9].

Not only does agriculture transfer land, but the changes in the use of this land also have great implications for microbial diversity. Conventional agricultural practices, such as over-application of chemical fertilizers and pesticides, known as monoculture and over-tillage, have been shown to cause loss of soil microbial populations to a large extent, resulting in degradation of soil fertility with time. It is the disruption of soil's natural structure, degradation of organic materials, and harm to beneficial microorganisms that resulted in a decline in soil fertility in the long run. On the other hand, sustainable agricultural practices such as organic farming practices, reduced tillage, and crop rotation increase microbial diversity and improve the habitat conditions for these organisms. For instance, organic practices attain an organic soil-matter buildup, thus excluding most of the synthetic chemicals in farming and creating a thriving community of microorganisms. [56].

One of the major functions of soil microorganisms consists of nutrient cycling. The soil microbes help in the cycling of quite essential nutrients available to plants like nitrogen, phosphorus, and potassium. The very important activity was nitrogen fixation, which was carried out by certain bacteria such as *Rhizobium* and *Azotobacter* in which they changed atmospheric nitrogen to forms that can be absorbed by plants. Similarly, phosphorus-solubilizing bacteria made phosphorus available to plants by mobilizing it from insoluble forms in soil. Potassium mobilization by soil microorganisms enhanced the availability of potassium which is another very essential nutrient for the growth of plants. All these activities are important for maintaining soil fertility and, hence, making nutrients available to plants for optimal growth [57].

Apart from the nutrient cycling envisioned by microorganisms, these organisms contribute substantially to the process of organic matter decomposition and humus formation. Besides, there are favorable conditions for the decay by bacteria, fungi, and actinomycetes of plant residues, animal remains, and other organics. This organic matter is then reduced from a complicated form to a simpler constituent, which will be integrated into the soil to produce the end product in the organometal association as one component of soil organic matter. It thus strengthens the soil structure, increases water retention, and boasts of holding nutrients in soils. This will create an environment in soils that will be sufficient for efficient decomposition and formation of stable humus, which boosts the productivity of soil towards accommodating the growth of plants. [57].

Soil structure as well as its aggregation processes are partly the work of microorganisms. Soil aggregates can be defined as groupings of soil particles that are held by organic matter and microorganisms. Binding agents like polysaccharides play a role in exuding microbes with stable aggregates. The soil structure should be built in such a way that it can enhance infiltration, root penetration, and gas exchange, which are all vital for the good health of plants. Moreover, soil aggregates hold particles together, preventing soil erosion and compaction, which in turn promote soil health and fertility [56].

Microorganisms also have an important role in regulating the soil pH. Some soil microbes can synthesize acids or alkaline products, thereby changing the pH of the soil. For example, certain bacteria produce organic acids that alkalify soil from iron and aluminum, making it more available to plants. Other microbes raise the soil pH and thus allow it to be suitable for other crops. Microorganisms regulate soil pH, thereby optimizing nutrient availability to plants and improving soil fertility [57].

The diversity of microbes has a considerable effect on plant growth through symbiotic relations. One such notable example is the relationship established between leguminous plants and nitrogen-fixing rhizobia. Rhizobia grows in root nodules, where they fix free atmospheric nitrogen and supply it to the plant in exchange for carbon. Mycorrhizae are other

organisms that form an endophytic symbiosis with plant roots, extending root surfaces and improving nutrient uptake, especially phosphorus. This relationship helps promote plant growth and contributes to soil fertility improvement by increasing the cycling and availability of nutrients. [57].

Another group of microorganisms that directly promote plant health is Plant growth-promoting rhizobacteria (PGPR). They promote plant growth through nutrient uptake enhancement, plant hormone production, and pathogen suppression. PGPR can also serve as biofertilizers for higher crop yields and fertility enhancement. Moreover, within plant tissues, endophytic microorganisms live without inflicting harm. Their function includes plant growth enhancement and stress resistance. They empower plants against environmental challenges such as drought, salinity, and diseases [56]. The modern techniques for assessing microbial diversity, such as molecular methods like 16S-rRNA sequencing, metagenomics, and metatranscriptomics, have expanded the already vast horizons of studying soil microbial communities. These techniques can reveal unknown microbes as well as provide clues about the functional roles of different microbial groups. By analyzing the genetic material of soil microorganisms, researchers will find a wider understanding of how microbial diversity affects soil health and fertility.

Conclusion:

Microbial diversity has a vital role to play in the sustenance and enhancement of soil fertility. Soil microorganisms participate in nutrient cycling, decomposition of organic matter, soil aggregation, and pH regulation in soils. Microbial diversity is integral to sustainable agricultural practices such as organic farming, microbial inoculants, and composting that strive to improve soil health and fertility. Through ongoing research in soil microbiology, we understand more of the intricate interactions involving soil microbes, plants, and the environment. This will in turn inform measures that will bring about improved and sustainable approaches to soil fertility management, particularly in the face of the broader challenges of climate change and population increase.

Acknowledgment:

We highly acknowledge The Islamia University of Bahawalpur and the University of Okara who provided us with the platform labs, and internet sources to search for the article.

Author contribution:

The authors confirm their contribution to the paper: study conception and design of Asiya Akbar, Data Collection by Dr. Saman Zahra, Analysis and interpretation of the result by Qasid Hussain, Draft, and manuscript preparation, Usman Waheed and, Muhammad Rehman. All authors reviewed the results and approved the final version of the manuscript.

Data Availability:

All the procedure is performed in the Lab and related data is collected from the authentic net resources.

Funding:

No funding was granted from any source.

REFERENCES

1. Datsko, O., et al., Increasing soil fertility as a factor in the sustainability of agriculture and resilience to climate change. 2024.
2. Gunal, H., et al., Soil fertility in arid lands: strategies for sustainable management and fertilization, in Sustainable Agriculture under Drought Stress. 2025, Elsevier. p. 61-69.
3. Patel, S., et al., SOIL FERTILITY AND NUTRIENT MANAGEMENT.
4. Gupta, A., et al., Linking soil microbial diversity to modern agriculture practices: a review. International Journal of Environmental Research and Public Health, 2022. 19(5): p. 3141.
5. Rao, D., Microbial diversity, soil health and sustainability. Journal of the Indian Society of Soil Science, 2007. 55(4): p. 392-403.
6. Nadarajah, K., Soil health: the contribution of microflora and microfauna. Mycorrhizosphere and Pedogenesis, 2019: p. 383-400.

7. Ogunseitan, O., Microbial diversity: form and function in prokaryotes. 2008: John Wiley & Sons.
8. Nannipieri, P., et al., Microbial diversity and soil functions. *European journal of soil science*, 2017. **68**(1): p. 12-26.
9. Kennedy, A.C. and K. Smith, Soil microbial diversity and the sustainability of agricultural soils. *Plant and soil*, 1995. **170**: pp. 75-86.
10. Akbar, A., et al., The Role of Sustainable Farming Practices in Enhancing Soil Fertility and Crop Yield. *Dialogue Social Science Review (DSSR)*, 2024. **2**(5): p. 137-147.
11. Pandey, S.N., M. Abid, and M.M. Abid Ali Khan, Diversity, functions, and stress responses of soil microorganisms. *Plant Microbiome: Stress Response*, 2018: p. 1-19.
12. THORN, R.G. and M.D. LYNCH, Fungi, and eukaryotic algae, in *Soil microbiology, ecology, and biochemistry*. 2007, Elsevier. p. 145-162.
13. Dighton, J., *Fungi in ecosystem processes*. 2018: CRC Press.
14. Sharma, M., P. Dangi, and M. Choudhary, Actinomycetes: source, identification, and their applications. *International Journal of Current Microbiology and Applied Sciences*, 2014. **3**(2): p. 801-832.
15. Provasoli, L., Nutrition, and ecology of protozoa and algae. *Annual review of microbiology*, 1958. **12**(1): p. 279-308.
16. Schlichting Jr, H., The importance of airborne algae and protozoa. *Journal of the Air Pollution Control Association*, 1969. **19**(12): p. 946-951.
17. Zhou, J., et al., Spatial and resource factors influencing high microbial diversity in soil. *Applied and Environmental Microbiology*, 2002. **68**(1): p. 326-334.
18. Zifcakova, L., Factors affecting soil microbial processes. *Carbon and Nitrogen Cycling in Soil*, 2020: p. 439-461.
19. Mohamed, H.I., et al., Role of microorganisms in managing soil fertility and plant nutrition in sustainable agriculture. *Plant growth-promoting microbes for sustainable biotic and abiotic stress management*, 2021: p. 93-114.
20. Lawlor, D.W., G. Lemaire, and F. Gastal, Nitrogen, plant growth, and crop yield. *Plant nitrogen*, 2001: p. 343-367.
21. Zhu, J., M. Li, and M. Whelan, Phosphorus activators contribute to legacy phosphorus availability in agricultural soils: A review. *Science of the Total Environment*, 2018. **612**: p. 522-537.
22. Malvi, U.R., Interaction of micronutrients with major nutrients with special reference to potassium. *Karnataka Journal of Agricultural Sciences*, 2011. **24**(1).
23. Condron, L., et al., The role of microbial communities in the formation and decomposition of soil organic matter. *Soil microbiology and sustainable crop production*, 2010: p. 81-118.
24. Masciandaro, G., et al., Organic matter-microorganism-plant in soil bioremediation: a synergic approach. *Reviews in Environmental Science and Bio/Technology*, 2013. **12**: p. 399-419.
25. Allison, F.E., *Soil organic matter and its role in crop production*. 1973: Elsevier.
26. Naz, M., et al., The soil pH and heavy metals revealed their impact on soil microbial community. *Journal of Environmental Management*, 2022. **321**: p. 115770.
27. Alexander, M., Effects of acidity on microorganisms and microbial processes in soil, in *Effects of acid precipitation on terrestrial ecosystems*. 1980, Springer. p. 363-374.
28. Wang, C.-y., et al., Soil pH is the primary factor driving the distribution and function of microorganisms in farmland soils in northeastern China. *Annals of Microbiology*, 2019. **69**(13): p. 1461-1473.
29. Das, P.P., et al., Plant-soil-microbes: A tripartite interaction for nutrient acquisition and better plant growth for sustainable agricultural practices. *Environmental Research*, 2022. **214**: p. 113821.
30. Martin, F.M., S. Uroz, and D.G. Barker, Ancestral alliances: plant mutualistic symbioses with fungi and bacteria. *Science*, 2017. **356**(6340): p. eaad4501.

31. Soto, M.J., et al., Mutualism versus pathogenesis: the give-and-take in plant-bacteria interactions. *Cellular microbiology*, 2009. **11**(3): p. 381-388.
32. Selosse, M.-A., E. Baudoin, and P. Vandenkoornhuys, Symbiotic microorganisms, a key for ecological success and protection of plants. *Comptes rendus biologies*, 2004. **327**(7): p. 639-648.
33. Santoyo, G., et al., Rhizosphere colonization determinants by plant growth-promoting rhizobacteria (PGPR). *Biology*, 2021. **10**(6): p. 475.
34. Goswami, D., J.N. Thakker, and P.C. Dhandhukia, Portraying mechanics of plant growth promoting rhizobacteria (PGPR): a review. *Cogent Food & Agriculture*, 2016. **2**(1): p. 1127500.
35. Saharan, B. and V. Nehra, Plant growth promoting rhizobacteria: a critical review. *Life Sci Med Res*, 2011. **21**(1): p. 30.
36. Islam, W., et al., Role of environmental factors in shaping the soil microbiome. *Environmental Science and Pollution Research*, 2020. **27**: p. 41225-41247.
37. Zheng, Q., et al., Soil multifunctionality is affected by the soil environment and by microbial community composition and diversity. *Soil Biology and Biochemistry*, 2019. **136**: p. 107521.
38. Ma, L., et al., Environmental factors and microbial diversity and abundance jointly regulate soil nitrogen and carbon biogeochemical processes in Tibetan wetlands. *Environmental Science & Technology*, 2020. **54**(6): p. 3267-3277.
39. Andrew, D.R., et al., Abiotic factors shape microbial diversity in Sonoran Desert soils. *Applied and environmental microbiology*, 2012. **78**(21): p. 7527-7537.
40. Peng, S.-l., et al., Environmental drivers of soil microbial activity and diversity along an elevational gradient. *Journal of Mountain Science*, 2022. **19**(5): p. 1336-1347.
41. Xia, Q., T. Ruffy, and W. Shi, Soil microbial diversity and composition: Links to soil texture and associated properties. *Soil Biology and Biochemistry*, 2020. **149**: p. 107953.
42. Sarethy, I.P., S. Pan, and M.K. Danquah, Modern taxonomy for microbial diversity. *Biodiversity-the dynamic balance of the planet*, 2014.
43. Hill, G., et al., Methods for assessing the composition and diversity of soil microbial communities. *Applied soil ecology*, 2000. **15**(1): p. 25-36.
44. Bing-Ru, L., et al., A review of methods for studying microbial diversity in soils. *Pedosphere*, 2006. **16**(1): p. 18-24.
45. Fukuda, K., et al., Molecular approaches to studying microbial communities: targeting the 16S ribosomal RNA gene. *Journal of UOEH*, 2016. **38**(3): p. 223-232.
46. Benlloch, S., A.J. Martínez-Murcia, and F. Rodríguez-Valera, Sequencing of bacterial and archaeal 16S rRNA genes directly amplified from a hypersaline environment. *Systematic and applied microbiology*, 1995. **18**(4): p. 574-581.
47. Rheims, H., F. Rainey, and E. Stackebrandt, A molecular approach to search for diversity among bacteria in the environment. *Journal of Industrial Microbiology*, 1996. **17**: p. 159-169.
48. Gantner, S., et al., Novel primers for 16S rRNA-based archaeal community analyses in environmental samples. *Journal of Microbiological Methods*, 2011. **84**(1): p. 12-18.
49. Janssen, P.H., Identifying the dominant soil bacterial taxa in libraries of 16S rRNA and 16S rRNA genes. *Applied and environmental microbiology*, 2006. **72**(3): p. 1719-1728.
50. Gajda, A.M., et al., Effects of different soil management practices on soil properties and microbial diversity. *International Agrophysics*, 2018. **32**(1): p. 81.
51. Rao, D. and A. Patra, Soil microbial diversity and sustainable agriculture. *Journal of the Indian Society of Soil Science*, 2009. **57**(4): p. 513-530.
52. Sofu, A., et al., Soil microbial diversity and activity in a Mediterranean olive orchard

- using sustainable agricultural practices. *Soil Use and Management*, 2014. **30**(1): p. 160-167.
53. Wen-Hui, Z. and C. Zu-Cong, Effect of soil management practices and environmental factors on soil microbial diversity: a review. *Biodiversity Science*, 2004. **12**(4): p. 456.
54. Bertola, M., A. Ferrarini, and G. Visioli, Improvement of soil microbial diversity through sustainable agricultural practices and its evaluation by-omics approaches: A perspective for the environment, food quality, and human safety. *Microorganisms*, 2021. **9**(7): p. 1400.
55. Yang, T., K.H. Siddique, and K. Liu, Cropping systems in agriculture and their impact on soil health-A review. *Global Ecology and Conservation*, 2020. **23**: p. e01118.
56. Kibblewhite, M., K. Ritz, and M. Swift, Soil health in agricultural systems. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 2008. **363**(1492): p. 685-701.
57. Attiwill, P.M. and M.A. Adams, Nutrient cycling in forests. *New phytologist*, 1993. **124**(4): p. 561-582.

